

Preliminary Astrometric Results from the PS1 Demo Month and Routine Survey Operations

David G. Monet¹,
Pan-STARRS Project,
Pan-STARRS PS1 Consortium

ABSTRACT

Pan-STARRS PS1 has transitioned from its development phase to its operational phase. In March 2010, the Demo Month used a special cadence to allow data exploration to start. In May 2010, the Operation Phase commenced. Although there have been occasional breaks and revisions, routine data collection has been maintained. So far, more than 16,000 frames are available to the PS1 Consortium for analysis. Based on these first few frames, the provisional astrometric results show the error for a single measure of a well-exposed star to be about 15 to 20 milliarcsec.

¹ U. S. Naval Observatory Flagstaff Station; *dgm@nofs.navy.mil*

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE SEP 2010		2. REPORT TYPE		3. DATES COVERED 00-00-2010 to 00-00-2010	
4. TITLE AND SUBTITLE Preliminary Astrometric Results from the PS1 Demo Month and Routine Survey Operations			5a. CONTRACT NUMBER		
			5b. GRANT NUMBER		
			5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U. S. Naval Observatory, Flagstaff Station, AZ, 86001-8521			8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES 2010 Advanced Maui Optical and Space Surveillance Technologies Conference, 14-17 Sep, Maui, HI.					
14. ABSTRACT Pan&#8208;STARRS PS1 has transitioned from its development phase to its operational phase. In March 2010, the Demo Month used a special cadence to allow data exploration to start. In May 2010, the Operation Phase commenced. Although there have been occasional breaks and revisions routine data collection has been maintained. So far, more than 16,000 frames are available to the PS1 Consortium for analysis. Based on these first few frames, the provisional astrometric results show the error for a single measure of a well&#8208;exposed star to be about 15 to 20 milliarcsec.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 7	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

1) Survey Operations:

The Pan-STARRS PS1 survey began its Operational phase in May 2010. Since then, there have been minor interruptions but the survey has been taking large amounts of very high quality data. The cadence optimizes sky coverage in multiple survey colors (SDSS g -, r -, i -, z -, y -bands) at the expense of repeated visits to the same field. The current sky coverage map shows this. Using a pseudo-color representation of the filters, Figure 1 shows the current sky coverage based on 16,747 full-camera observations (60 Orthogonal Transfer CCD images each) transferred to the U.S. Naval Observatory Flagstaff Station (NOFS) so far. More frames have been taken, and this count does not include observations taken prior to the Demo Month.




Figure 1: PS1 sky coverage as of September 1, 2010.

2. Correlation With Stars In Other Catalogs:

The PS1 Image Processing Pipeline ² (IPP) correlates the images detected on each frame with the 2MASS catalog ³ to produce preliminary astrometric solutions. Using these coordinates, the NOFS astrometric pipeline correlates PS1 objects with other catalogs (USNO-B1.0, SDSS, Tycho-2, 2MASS, etc.) so that unique star identification numbers can be assigned across all catalogs. This correlation is done in two steps. The first step uses a 1.0 arcsec radius, and when completed the correlated PS1 observations and their reference catalog counterparts are removed from the lists. The second step uses a 10.0 arcsec radius. Figure 2 shows the number of correlations as a function of radius for the 345,264,598 PS1 objects with counterparts in other catalogs.




Figure 2. Number of correlations as a function of radius.

² <http://pan-starrs.ifa.hawaii.edu/public/project-status/project-status.html>

³ <http://irsa.ipac.caltech.edu/Missions/2mass.html>

Figure 2 is interpreted in the following manner. The initial steep decline is the astrometric error for each correlated pair, and the standard deviation for these pairings is about 0.3 arcsec. Whereas the 2MASS catalog error for a brighter star is believed to be in the range of 0.07 to 0.10 arcsec, the luminosity function has more stars at fainter magnitudes where the astrometric error is larger due to simple photon statistics. Also, many PS1 stars do not appear in the 2MASS catalog, and the astrometric error of the USNO-B1.0 catalog is believed to be in the range of 0.2 to 0.3 arcsec.

Hence, the astrometric error for the typical IPP coordinate is probably in the range of 0.1 to 0.2 arcsec depending on brightness. The break at 1.0 arcsec arises from the two-part nature of the search. The change of slope and structure between about 1 and 5 arcsec is interpreted as the detection of true stellar proper motions. The change of slope and slow rise in the number of correlations for radii larger than 5 arcsec are interpreted as effects of spurious correlations: there should be a very few true proper motion stars with such motions. These stars would be quite exciting to find, but are very difficult to confirm.

3) Astrometric Solutions:

The NOFS astrometric pipeline was designed to deal with large quantities of high quality data, but it faces many serious challenges. The PS1 cadence includes dithers of the telescope boresight and deliberate rotations of the camera with respect to the telescope in addition to the natural rotation associated with an alt-azimuth telescope design.

Hence, the HEALPix⁴ tessellation scheme was adopted, and *side*=64 was used for the data storage strategy and *side*=256 was used for the individual astrometric solutions. A great deal of bookkeeping is needed to identify which OTCCD of which observation measured stars in a specific HEALPix. In many cases, each HEALPix is spread over several OTCCDs. A separate astrometric solution is needed for each OTCCD for each observation for each HEALPix. So far, there are multi-epoch PS1 observations in 335,231 *side*=256 HEALPix.

Interpretation of the ensemble of HEALPix solutions is made more complicated by several factors. Perhaps the most important is the current inability of the NOFS pipeline to compute and incorporate the weights appropriate for a particular observation. Effects such as seeing, sky background, focus, and wind should be incorporated into the weights, but at the moment all observations are assigned unit weight. Another significant problem is the very preliminary nature of the algorithm designed to sense and remove statistical outliers. There are many reasons why the measured position could be in error, but without a reliable master catalog it is quite difficult to sense these in the ensemble of measures. Also, the effects of differential color refraction (DCR) should displace the observed positions as a function of star color and the circumstances of the observation. Again, the lack of a master catalog makes it difficult to fit such terms. With these caveats, statistics from the analysis of about 9,700 well-observed HEALPix are shown in Figure 3.

⁴ <http://healpix.jpl.nasa.gov>




Figure 3: Astrometric error distributions in (in milliarcsec) for single color solutions in $side=256$ HEALPix with PS1 observational epochs covering more than 30 days. The red lines are from the ξ (RA) solutions and the blue are from η (Dec) solutions.

In spite of all of the difficulties mentioned above, it seems reasonable to conclude that under good circumstances (whatever this really means) that a single measure of a well exposed stellar image has an astrometric error between 15 and 20 milliarcsec (mas) in each coordinate. The long tails shown in Figure 3 are interpreted as being the results of the inadequate trimming of statistical outliers and the incomplete astrometric model. The solutions shown in Figure 3 include only an Affine transformation between the OTCCD (column,row) and tangent plane coordinates (ξ,η). The size of a *side*=256 HEALPix is about 14 arcmin on a side, but there may be significant focal plane curvature over this size, particularly near the edge of the field of view. The impacts of DCR are still unknown.

4) Conclusions:

The results presented here are quite preliminary, and reflect the mis-match between the task to be done and the level of effort available. However, the results presented in Figure 3 offer some hope that the astrometric error of a single observation is between 15 and 20 mas. This is about a factor of 2 worse than the pre-survey predictions, but much work remains to be done. The most important areas for software development are weights computed from the metadata , an improved algorithm for sensing and removing statistical outliers, and the removal of the effects of DCR.

It is DGM's pleasure to thank the members of the PS1 Project and Consortium teams. This is an amazing project!